


U.S. Gulf of Mexico Sea Turtle Habitat Areas Loggerhead



Five species of sea turtles are found in Gulf waters, all of which are either threatened or endangered. Prime nesting season for sea turtles is May through October. Green sea turtles consume seagrasses and algae, loggerheads forage benthically and consume crustaceans and fish, leatherbacks eat primarily jellyfish, and Kemp's ridleys are normally bottom-feeders, consuming crabs and other benthic fauna. USGS is currently conducting sea turtle sampling on juvenile greens and adult loggerheads in Everglades National Park and on several life stages of loggerheads, greens, and hawksbills in Dry Tortugas National Park. The work in the Dry Tortugas is supported by both the Coral Reef Ecosystem Studies Project (CREST) and the Priority Ecosystem Studies (PES) Program. The map depicts locations of current sea turtle sampling projects (stars) and polygons showing results of previous sea turtle tracking projects by species in relation to the BP Deepwater Horizon oil wellhead. Also shown are results from previous aerial surveys to document sea turtle presence in the Gulf and Atlantic coastal areas.

Polygons may include locations for more than one turtle, and nesting sites are often used by several species of sea turtles.

This map is preliminary, for informational purposes only.

Data Sources:

- 1. Fritts, T.H., Hoffman, W. and M.A. McGehee. 1983. The distribution and abundance of marine turtles in the Gulf of Mexico and nearby Atlantic waters. Journal of Herpetology, 17:4 327-344.
- 2. Girard, C. Tucker A.D. and B. Calmettes. 2009. Post-nesting migrations of loggerhead sea turtles in the Gulf of Mexico: dispersal in highly dynamic conditions. Marine Biology 156: 1827-1839.
- 3. Hoffman, W. and T. H. Fritts. 1982. Sea turtle distribution along the boundary of the Gulf Stream current off eastern Florida. Herpetologica, 38:3 405-409.
- 4. Inwater Research Group. 2005. Key West Natoinal Wildlife Refuge Sea Turtle Project, 2005 Annual Report. Report submitted to NMFS, Silver Spring, MD.
- 5. Inwater Research Group. 2006. Population assessment of marine turtles in Lake Worth Lagoon, FL. Report submitted to Palm Beach County Dept of Environmental Resources Management, West Palm Beach, FL.
- 6. Inwater Research Group. 2005. Abundance and distribution of marine turtles within nearshore hardbottom and associated habitats. Report prepared for Florida Fish abd Wildlife Conservation Commission, St Petersburg, FL.
- 7. Norris, J.C., Evens, W.E. and S. Rankin. 2000. An Acoustic Survey of Cetaceans in the Northern Gulf of Mexico. Chapter 5 in: Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: distribution, abundance and habitat associations. Volume II: Technical Report. R.W. Davis, W.E. Evens and B. Wursig editors. U.S. Geological Survey, Biological Resources Division, by The GulfCet Program, Department of Marine Biology, Texas A&M University at Galveston.
- 8. Renaud, M.L. and J.A. Carpenter. 1994. Movements and submergence patterns of loggerhead turtles (Caretta caretta) in the Gulf of Mexico determined through satellite telemetry. Bulletin of Marine Science, 55:11-15.
- 9. Schmahl, G.P. 2003. Managing Marine Areas: Sucess stories. Flower Garden Bnaks National MArine Sanctuary. Presentation June 12, 2003. NOAA.
- 10. Witzell, W.N. and J.R. Schmid. 2004. Immature sea turtles in Gullivan Bay, Ten Thousand Islands, Southwest Florida. Gulf of Mexico Science, 1: 54-61.

11. USGS□12. NOAA 13. USGS CREST Project

14. Mote Marine Lab, Tony Tucker

15. Conservacy of SW Florida, David Addison

Map 1 of 5

16. USGS, Ray Carthy and Kristen Hart Personnal Comm.